

# Fact Sheet

## Aberdeen Selection of Laurel Willow

**S**pecies: *Salix pentandra*  
Common Name: Laurel Willow  
Plant Symbol: SAPE  
Accession Number: 9005049

**Source:** The Natural Resources Conservation Service, Plant Materials Center at Aberdeen, Idaho received the original plant material from the Plant Materials Center in Rose Lake, Michigan. It was collected in the upper Midwest from naturalized stands.

**Native Site Information:** This plant is native to central and western Europe where it has been cultivated for many centuries. Another common name is Bay Willow. Both common names refer to the similarity of the leaf to that of the Bay Laurel tree.

**Method of Selection:** Selected from a collection of potential windbreak plants assembled and evaluated at the Plant Materials Center from 1981 through 1996. Laurel willow, 9005049, has been tested at Off-Center locations near Mountain Home, Idaho, and Winnemucca, Nevada. This accession was selected for its beauty, hardiness, appropriate growth form for windbreaks, and natural range of adaptability. This accession shows great promise for use throughout the Intermountain West when used in irrigated windbreaks and landscaping.



LAUREL WILLOW

**Description:** Introduced from Europe during the past three centuries, Laurel willow has lustrous dark green foliage. The dark green, shiny leaves are 2-5 inches long and very attractive. The bark on the fine twigs and the stems is green to yellow green and gray on the older trunks. It requires deep moist soils or supplemental irrigation. It is a medium to tall shrub with a dense round top, symmetrical crown and multiple trunks. Laurel willow grows to heights of 20-40 feet (6-12 meters) with average canopy coverage of 15-25 feet (4-8 meters).



**Insect and Disease Problems:** No problems were observed with insects or diseases which affected the survival or appearance of this plant.

**Anticipated Use:** The potential uses of Laurel willow are wind erosion control, diversity and beautification. Its fibrous root system and wide canopy cover make it an excellent plant for soil stabilization. The moderately dense stem and leaf pattern make this an excellent plant for windbreaks. It is recommended for use in interior rows of multiple-row windbreaks, in single row or twin-row windbreaks where an evergreen is not needed or desired, for landscaping, and to provide nesting and roosting habitat for birds..

**Environmental Considerations:** Laurel willow is not intended for use in natural wetlands or riparian areas. Since it is an introduced plant from Europe, it is not an appropriate component in native plant communities. In riparian areas, stems cut by beaver and broken stems may spread this plant to areas beyond the landowner's control. The material being released is all from female plants.

**Potential Area of Adaptation:** The range of adaptation is very broad because the plant is expected to be used under managed conditions where annual rainfall is high (at least 22 inches per year) or where irrigation water is made available. It is tolerant of very cold weather and adapted for use in windbreaks and landscaping in all of the Intermountain West. We expect it will also perform well in the northern Great Plains and upper Midwest.

**Potential Soil Adaptation:** Deep or moderately deep loams, sandy loams, gravelly loams, well-drained to somewhat poorly drained soils.

**Harvesting and Planting:** This plant is reproduced using cuttings. Cuttings should be made during the period from late Fall to

early Spring when the plants are dormant and have no leaves. Limbs from .5 inch to two inches in diameter make excellent cuttings. All of the small side branches should be removed, as well as the upper part that is less than .25 inches in diameter. Cuttings can be from 3 to 6 feet long. Cuttings can be stored under refrigeration until Spring, or they can be planted immediately. The key to successful planting is to get the bottom of cutting as deep as possible. A good rule of thumb is to place at least 2/3 of the cutting in the ground and more if possible. Where irrigation water is provided, planting deep (and watering deep) are essential to long term survival. If the cuttings have been stored for a long period prior to planting, soaking the lower ends (all that will be in the ground) in water 1-4 days will speed up the rooting process. The use of root promoting hormones is not required. The scarring or cutting of the planted portion is not recommended and may hurt the plant.

**Parent Plant Maintenance:** Parent plants will be maintained at:  
USDA-NRCS; Aberdeen PMC; P.O. Box 296;  
1691A. S. 2700 W.; Aberdeen, ID 83210.  
PH: (208) 397-4133

Cuttings from Laurel willow will be made available to nurseries at no cost. Nurseries may establish their own parent plants for long-term production of cuttings.

January 1998



The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET CENTER at (202) 720-2791.

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C., 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal opportunity employer.